

BOWEL PREPARATION INSTRUCTIONS FOR A COLONOSCOPY

PICOSALAX

It is important that you notify us well in advance prior to using this regime if:

- You suffer from constipation, have had part of your bowel removed, have had a poor bowel preparation in the past, or have not tolerated the prep in the past
- Have diabetes requiring insulin and have not been given instructions on what to do with your insulin
- *Are on Semaglutide (eg Ozempic, Saxenda, Wegovy)*
- Are on blood thinning medications other than aspirin (eg Warfarin, Coumadin, Clopidogrel, Plavix, Rivaroxaban, Xarelto, Apixaban, Eliquis, Dabigatran, Pradaxa, Ticlopidine, Ticlid, Ticagrelor, Brillinta, Iscover, Heparin, Clexane, Fragmin, Dipyridamole, Asasantin, Persantin, etc)

These are instructions on how to ensure that your colon is clean prior to your colonoscopy. It is very important for it to be clean so that the inside of your colon can be inspected thoroughly. If you do not follow the bowel preparation instructions and too much faecal material is still present then the procedure may need to be repeated.

Seven days before your procedure:

- Stop taking iron supplements
- You will need to obtain four sachets of Picosalax (or Picoprep or Picolax if Picosalax is not available) from your local pharmacist. They may need to order it in for you and hence why we suggest you purchase it well in advance of your procedure.

Three days before your procedure:

- Stop taking any drugs that may make you constipated (eg Codeine, Panadeine Forte, Tramadol, Oxycodone etc). If you cannot do without these drugs please contact the rooms for special advice
- You must now begin eating only a low residue diet. Please see page 4 for examples of foods you may eat.
- It is really important that you do not eat foods that are not on the list, especially those with nuts and seeds, as these are hard to digest, and obscure the views at the time of the colonoscopy, which can sometimes mean the procedure needs to be repeated. Stop eating all fruits, muesli, bran, high fibre cereals, nuts and anything with seeds in it (eg multigrain bread)

FOR A MORNING PROCEDURE (Before 1pm)

The Day before your procedure

1. You may eat a low residue breakfast and lunch before midday (See page 4 for allowed foods).
2. After midday you may only drink clear fluids (See page 3 for allowed clear fluids). No more food is now allowed until after the procedure.
3. At 1pm take the first sachet of Picosalax, then drink 500ml of clear fluids.

DR. AMMAR NAGREE
(Principal)

Gastroenterologist
MBBS (Syd Uni), FRACP
Provider number 206667AT

DR. KATHERINE BROUGHTON

Colorectal Surgeon / Endoscopist
MBChB, BHB, FRACS
Provider number 509951CF

DR. ADRIAN MCELHOLM

Gastroenterologist
MMedSc FRACP (Glasg) FRACP MD
Provider number 424241H

DR. MUKESH NASA

Gastroenterologist
MBBS MD DM FRACP
Provider number 549228BJ

DR. SIVA PULUSU

Gastroenterologist
MBBS FRACP
Provider number 5253445B

4. At 5pm take the second sachet of Picosalax, then drink 500ml of clear fluids.
5. At 9pm take the third sachet of Picosalax, then drink 500ml of clear fluids.

The day of your procedure

1. Four hours before your admission time take the fourth sachet of Picosalax, then drink 500mls of clear fluids.
2. In the two hours before your admission time you must completely fast from all food and liquids. If you have anything to eat or drink or if you chew gum in this two hour pre procedure period the procedure may need to be cancelled.

FOR AN AFTERNOON PROCEDURE (1pm or after)

The Day before your procedure

1. You may eat a low residue breakfast and lunch and light dinner before 5pm. (See page 4 for allowed foods).
2. After 5pm you may only drink clear fluids (See page 5 for allowed clear fluids). No more food is now allowed until after the procedure.
3. At 5pm take the first sachet of Picosalax, then drink 500ml of clear fluids.
4. At 9pm take the second sachet of Picosalax, then drink 500ml of clear fluids.

The day of your procedure

1. At 6am take the third sachet of Picosalax, then drink 500mls of clear fluids.
2. Four hours before your admission time take the fourth sachet of Picosalax, then drink 500mls of clear fluids.
3. In the two hours before your admission time you must completely fast from all food and liquids. If you have anything to eat or drink or if you chew gum in this two hour pre procedure period the procedure may need to be cancelled.

SPECIAL NOTES:

- If you develop nausea with drinking the preparation then you may take your time to drink it slower. It is quite sweet so you can mix it with another clear fluid to neutralize the taste (eg black tea, black coffee, strained lemon juice etc)
- If at any stage you vomit the preparation, or you have any other concerns regarding preparation please telephone my consulting rooms (9528 1192). If you need advice after hours then, depending on which hospital your procedure has been booked at, contact Waikiki Private Hospital 9550 0222 (ask to speak to one of the ward nurses). If you are unable to contact them then phone your after-hours GP, the Nurse Direct Line (1800022222) or your local emergency department.
- Once your laxatives begin to take effect you will lose a lot of fluid and it is critical that you maintain your hydration. Do not over exert yourself once you have begun the prep just in case your blood pressure begins to fall as a result of dehydration. Feeling faint can occasionally occur and should be treated by drinking more fluids and resting.
- You should take any medications (other than iron and constipating medications) that you would normally take up until the two hours before your admission time. Any medications that would normally be taken in the two hours before your admission time do not need to be taken.
- If you are diabetic then you should take your diabetic tablets as normal. If you are requiring insulin then you should have already contacted my rooms in advance to ask for special advice. If you haven't done so then immediately call one of the numbers listed below.
- You may use a barrier cream such as castor oil on your bottom, if required, to prevent soreness.
- Some intestinal cramping is normal.
- In the two hours before your procedure when you are fasting from all food and liquids, chewing gum is also not allowed.
- Your stools should be clear or yellow. If they are not then please come to the hospital 30 minutes before your appointment time, as you may require further laxatives.

Clear fluids include the following:

- Water
- Black tea
- Black coffee
- Cordial – orange or lemon **NO red, green, purple or blue**
- Fruit juice – pulp free orange or clear apple
- Clear Soup - strain bits out before drinking
- Yellow or orange jelly
- Sports Drinks eg Gatorade or Powerade. **NO red, green, purple or blue**
- Beef tea eg Bovril, Bonox
- Soup cubes eg stock cube in hot water

Suggested 'low residue' menu plan only

The menu plan is a suggestion only and is not intended for long term use, as it contains fewer vegetables and fruits than recommended in a balanced diet.

BREAKFAST	LUNCH/DINNER	SNACKS
Juice – apple, cranberry, pear, white grape	Clear soups, broths, chicken noodle soup	Cheese and plain crackers
Plain yoghurt (no fruit or nut topping)	Fish, veal, lean lamb, lean pork, chops, ham, chicken, turkey	Rice cakes, protein bars (no nuts)
Small bowl Cornflakes or Rice bubbles	Potato or pumpkin (skinless), baked, steamed, mashed, potato salad	Ham quiche
2 eggs poached, boiled or scrambled	Pasta carbonara or macaroni	Parmesan cheese sticks
Bacon, sausage	Cheese (no tomato paste or bottled sauces)	Wedges with sour cream
Hash browns	Sandwiches - white bread/roll	Cake based on white flour
1 slice white toast	Panini: chicken, ham, (no mustard), tuna, turkey, pastrami, corned beef, cheese, egg	Biscuits based on white flour
French toast, cinnamon sugar	Baked rice custard, chocolate, mousse	Chocolate biscuits - no nuts
Vegemite, honey	Plain ice cream	Cheesecake - no whole meal crust
Croissants - plain or ham and cheese	Water, soft drink, tea, coffee	Scones with honey
Low fibre muffin - no fruits or nuts		Pikelets with lemon & sugar or honey
Crumpets and honey		Waffles with syrup
Pancakes with syrup		Vanilla slice (no passionfruit)
Chocolate brioche		Doughnut, danish, custard tart, meringue kisses
Tea, coffee, water		Milkshakes, protein shakes
		Clear sports drinks, tea, iced tea, coffee, hot chocolate
		Plain/chocolate Sustagen
		Ensure, Digestelact, water